

ON THE DEPTH OF BLOW-UP RINGS OF IDEALS OF MINIMAL MIXED MULTIPLICITY

CLARE D'CRUZ

ABSTRACT. We show that if $(R, \mathfrak{m})$ is a Cohen-Macaulay local ring and I is an ideal of minimal mixed multiplicity, then $\text{depth } G(I) \geq d - 1$ implies that $\text{depth } F(I) \geq d - 1$. We use this to show that if I is a contracted ideal in a two dimensional regular local ring then $\text{depth } R[It] - 1 = \text{depth } G(I) = \text{depth } F(I)$. We also give an infinite class of ideals where $R[It]$ is Cohen-Macaulay but $F(I)$ is not.

1. INTRODUCTION

Throughout this paper we will assume that $(R, \mathfrak{m})$ is a local ring of dimension d with infinite residue field. Contracted ideals in a two dimensional regular local ring have been studied by Zariski [ZS]. A special class of contracted ideals are complete ideals. Zariski's work reveals that complete ideals satisfy several interesting properties. We would like to know if some of these results hold good for contracted ideals.

Associated to any ideal I in local ring $(R, \mathfrak{m})$ we have the blowup rings, namely the the Rees ring $R(I) := R[It] = \bigoplus_{n \geq 0} I^n t^n$, the associated graded ring $G(I) := R(I) \otimes_R R/I = \bigoplus_{n \geq 0} I^n / I^{n+1}$ and the fiber cone $F(I) := \bigoplus_{n \geq 0} I^n / \mathfrak{m} I^n$. It is well known that all the blowup rings associated to a complete ideal in a two dimensional regular local ring are Cohen-Macaulay. One would naturally address the same question for contracted ideals. An interesting result which plays an important role in the Cohen-Macaulayness of blowup rings is the well known result of Goto and Shimoda which gives a necessary and sufficient condition for the Rees ring to be Cohen-Macaulay [GS]. From their result it follows that if $(R, \mathfrak{m})$ is a Cohen-Macaulay local ring, I is an $\mathfrak{m}$ -primary ideal, and the Rees ring is Cohen-Macaulay, then so is the associated graded ring. By using the Briancon-Skoda theorem, it was shown that for a regular local ring $(R, \mathfrak{m})$, the ring $R(I)$ is Cohen-Macaulay if and only if $G(I)$ is Cohen-Macaulay for any $\mathfrak{m}$ -primary ideal [AH]. We have extended this result of Huneke to relate the depth of the Rees ring and the associated graded ring of and $\mathfrak{m}$ -primary ideal in a regular local ring of positive dimension (see Theorem 2.2).

Motivated by these results, a natural question to ask is whether the depth of the fiber cone is related with the depth of the other blowup rings. For complete ideals in a regular local ring, all the blowup rings are Cohen-Macaulay. In [J], it was shown that if I is a lex-segment ideal in a two-dimensional polynomial ring, then $\text{depth}(R(I)) = \text{depth}(G(I)) = \text{depth}(F(I))$. In this paper we show that this result holds true for any $\mathfrak{m}$ -primary contracted ideal in a two dimensional regular local ring (see Theorem 3.8).

Contracted ideals in a two dimensional regular local ring are precisely ideals of minimal mixed multiplicity (Lemma 3.3). For an $\mathfrak{m}$ -primary contracted ideal I in a two dimensional regular ring $(R, \mathfrak{m})$, $\text{depth}(R(I)) - 1 =$

$\text{depth}(G(I)) = \text{depth}(F(I))$ (see Theorem 3.8). Since $\mathfrak{m}$ -primary contracted ideals in a two dimensional regular local ring are ideals of minimal mixed multiplicity, we study ideals of minimal mixed multiplicity in a Cohen-Macaulay local ring of dimension at least two. For ideals of minimal mixed multiplicity we show that if $G(I)$ is Cohen-Macaulay, then $\text{depth}(F(I)) \geq d - 1$ (see Proposition 3.6).

Several examples of $G(I)$ being Cohen-Macaulay but $F(I)$ is not is known, but an example of the Rees ring being Cohen-Macaulay but $F(I)$ is not was not known. In Example 4.1 we produce an infinite class of complete ideals in a regular local ring of dimension three where the ideal Rees ring is Cohen-Macaulay but fiber cone is not. The next question of interest is: If $F(I)$ is Cohen-Macaulay, is $G(I)$ and $R(I)$ Cohen-Macaulay. This question has a positive answer when I is an ideal of minimal mixed multiplicity (see Theorem 3.4).

I am very grateful to J. K. Verma who motivated me into this question. I also thank I. Swanson for encouraging me to write this paper.

2. DEPTH OF $R(I)$, $G(I)$ AND $F(I)$

In this section we first prove Theorem 2.2 which relates the depth of the Rees ring and the associated graded ring of $\mathfrak{m}$ -primary ideals in a regular local ring. Our main result is for contracted ideals. We first show for and $\mathfrak{m}$ -primary ideal in a two dimensional regular local ring, Cohen-Macaulayness of $R(I)$ implies the Cohen-Macaulayness of $G(I)$ and $F(I)$. We show that if I is an $\mathfrak{m}$ -primary ideal in a two dimensional regular local ring then $G(I)$ is Cohen-Macaulay if and only if $\text{depth}(F(I))$ is Cohen-Macaulay.

Let I be an ideal in a regular local ring $(R, \mathfrak{m})$. The integral closure of an ideal, denoted by $\bar{I}$ is defined as $\bar{I} = \{x \in R \mid x^n + a_1 x^{n-1} + \cdots + a_n = 0; a_i \in I^i\}$. An ideal I is complete if $I = \bar{I}$.

Theorem 2.1. *Let I be a complete ideal in a two dimensional regular local ring. Then $R(I)$, $G(I)$ and $F(I)$ are all Cohen-Macaulay.*

Proof. A complete ideal has $r(I) = 1$. It is well known that $G(I)$ is Cohen-Macaulay. Hence $R(I)$ is also Cohen-Macaulay [GS]. Cohen-Macaulayness of $F(I)$ follows from [Sh]. $\square$

We now study the depth of contracted ideals in a two-dimensional regular local ring. An ideal I is contracted if there exists $S = R[m/x]$ such that $I = IS \cap R$. In a two-dimensional regular local ring an $\mathfrak{m}$ -primary complete ideal is always contracted [ZS]. We prove this result for any dimension d .

We begin with a more general result which relates the depth of the Rees ring and the associated graded ring.

Theorem 2.2. *Let I be an $\mathfrak{m}$ -primary ideal in a regular local ring. Then $\text{depth}(R(I)) = \text{depth}(G(I)) + 1$.*

Proof. Since R is a regular local ring the ring $R(I)$ is Cohen-Macaulay if and only if $G(I)$ is Cohen-Macaulay for any $\mathfrak{m}$ -primary ideal [AH]. If $G(I)$ is not Cohen-Macaulay for an $\mathfrak{m}$ -primary ideal I , then $\text{depth } R(I) = \text{depth } G(I) + 1$ [HM]. Hence, $\text{depth } R(I) = \text{depth } G(I) + 1$ for all $\mathfrak{m}$ -primary ideals in a regular local ring. $\square$

A natural question to ask: If the Rees ring is Cohen-Macaulay, then is the fiber cone also. We have a positive answer for $\mathfrak{m}$ -primary ideals in a two dimensional regular local ring.

Corollary 2.3. *Let $(R, \mathfrak{m})$ be a two-dimensional regular local ring. Let I be an $\mathfrak{m}$ -primary ideal of R . If $R(I)$ is Cohen-Macaulay, then $G(I)$ and $F(I)$ are Cohen-Macaulay.*

Proof. The Cohen-Macaulayness of $G(I)$ follows from the Goto-Shimoda theorem. In a two dimensional Cohen-Macaulay local ring, if the Rees ring $R(I)$ is Cohen-Macaulay, then the reduction number of the ideal I is one by the Goto-Shimoda theorem. This in turn implies that the fiber cone $F(I)$ is Cohen-Macaulay [Sh]. $\square$

Remark 2.4. (1) *The converse of Corollary 2.3 is not true for any arbitrary $\mathfrak{m}$ -primary ideal. See Example 4.2.*
 (2) *Examples where the associated graded ring is Cohen-Macaulay but the fiber cone is not are well known ([HKU, Example 2.8]) and vice versa.*

In this section we give a partial answer to the converse of Corollary 2.3.

Proposition 2.5. *Let $(R, \mathfrak{m})$ be a two dimensional regular local ring and let I be an $\mathfrak{m}$ -primary contracted ideal of R . Then $F(I)$ is Cohen-Macaulay if and only if $G(I)$ is Cohen-Macaulay.*

Proof. Since I is contracted we know that if $F(I)$ is Cohen-Macaulay, if and only if $r(I) \leq 1$ [DRV]. Also since I is an ideal in a regular local ring, $G(I)$ is Cohen-Macaulay if and only if $r(I) = 1$. Hence $F(I)$ is Cohen-Macaulay if and only if $G(I)$ is Cohen-Macaulay. $\square$

To prove the equality of depth of fiber cone and associated graded ring for contracted ideals, we need the concept of minimal mixed multiplicity.

3. IDEALS OF MINIMAL MIXED MULTIPLICITY AND THE COMPLEX $C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))$

In this section we prove that for contracted ideals in a two dimensional regular local ring, the fiber cone and the associated graded ring have the same depth. We first show that in a two dimensional regular local ring, complete ideals are precisely examples of ideals of minimal mixed multiplicity. We first show that contracted ideals in a two-dimensional regular local ring are ideals of minimal mixed multiplicity. Next we consider the depth of the associated graded ring and fiber cone of an ideal of minimal mixed multiplicity in and Cohen-Macaulay ring of dimension at least two.

We show that if I is an ideal of minimal mixed multiplicity in a Cohen-Macaulay local ring of dimension at least two and if the associated graded ring is Cohen-Macaulay then the fiber cone has almost maximal depth. We exploit the graded complex $C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))$, where $x \in \mathfrak{m}$ and $\mathbf{a}_{d-1} = a_1, \dots, a_{d-1} \in I$.

Let I be an $\mathfrak{m}$ -primary ideal in a local ring of dimension d . It was proved in [B], that for large values of r and s , $\ell(R/\mathfrak{m}^r I^s)$ can be written as a polynomial in r and s and can be written in the form

$$\sum_{i+j \leq d} e_{ij} \binom{r+i}{i} \binom{s+j}{j}$$

where e_{ij} are integers. When $i+j = d$, the integers we put $e_{ij} = e_j(\mathfrak{m}|I)$ for $j = 0, \dots, d$. In this case these are called the mixed multiplicities of $\mathfrak{m}$ and I .

Definition 3.1. [DRV, Definition 2.2] *We say that an ideal has minimal mixed multiplicity if $\mu(I) = e_{d-1}(\mathfrak{m}|I) + d - 1$.*

Definition 3.2. *A set of elements $x_1, \dots, x_d$ is called a joint reduction of a set of ideals $I_1, \dots, I_d$ if $x_i \in I_i$ for $i = 1, \dots, d$ and there exists a positive integer n so that*

$$\left[\sum_{j=1}^d x_j I_1 \cdots \hat{I}_j \cdots I_d \right] (I_1 \cdots I_d)^{n-1} = (I_1 \cdots I_d)^n.$$

Rees proved that if $R/\mathfrak{m}$ is infinite, then joint reductions exist. From a result of Rees, it follows that $e_j(\mathfrak{m}|I)$ is the multiplicity of any joint reduction of the multi-set of ideals consisting of j copies of I and $d-j$ copies of $\mathfrak{m}$. We shall denote such a multi-set of ideals by $(\mathfrak{m}^{[d-j]}|I^{[j]})$.

Lemma 3.3. *Let I be an $\mathfrak{m}$ -primary ideal in a two dimensional regular local ring $(R, \mathfrak{m})$. Then I is a contracted ideal if and only if I is an ideal of minimal mixed multiplicity.*

Proof. Every contracted ideal in a regular local ring $(R, \mathfrak{m})$ satisfies the property that $\mu(I) = 1 + o(I)$, where $o(I)$ is the $\mathfrak{m}$ -adic order of I and $o(I) = e(\mathfrak{m}|I)$ [V, Theorem 4.1].

Conversely, suppose I is an ideal of minimal mixed multiplicity, then there exists a joint reduction (x, a) of $(\mathfrak{m}, I)$ such that $\mathfrak{m}I = xI + a\mathfrak{m}$. By [AH], to show that I is contracted it is enough to show that $\mathfrak{m}I : (x) = I$.

Clearly, $I \subseteq \mathfrak{m}I : (x)$. Now suppose $\alpha \in \mathfrak{m}I : (x)$. Then $\alpha x \in \mathfrak{m}I = xI + a\mathfrak{m}$. Hence $\alpha x = xi + a\beta$ for some $i \in I$ and $\beta \in \mathfrak{m}$. Therefore $x(\alpha - i) = a\beta$. Since (x, a) is a regular sequence in R , $\alpha - i = at$ for some $t \in R$. Hence $\alpha = i + at \in I$ since $i, a \in I$. $\square$

We first prove some results on ideals of minimal mixed multiplicity.

Theorem 3.4. *Let I be an ideal of minimal mixed multiplicity in Cohen-Macaulay local ring of dimension $d \geq 2$. If $F(I)$ is Cohen-Macaulay, then $G(I)$ and $R(I)$ are Cohen-Macaulay.*

Proof. From [DRV, Corollary 2.5], $F(I)$ is Cohen-Macaulay if and only if $r(I) \leq 1$. If $r(I) \leq 1$, then both $G(I)$ is Cohen-Macaulay and hence $R(I)$ is Cohen-Macaulay. $\square$

The converse of Theorem 3.4 need not always hold true, (see Proposition 3.6, Example 4.1). To prove this we consider a complex $C(\mathfrak{a}_{d-1}, \mathcal{F}, (1, n))$ which will link the depth of the associated graded ring with the depth of the fiber cone.

Let $\mathcal{F} = \{\mathfrak{m}I^n\}_{n \geq 0}$ be the filtration and let $x \in \mathfrak{m}$ and $\mathbf{a}_{d-1} \in I$. The complex $C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n))$, $C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n))$ $C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))$ can be constructed via the mapping cylinder. For details on these complexes we request the reader to refer to [HM] and [DG]. We have the short exact sequence:

$$0 \longrightarrow C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n)) \longrightarrow C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n)) \longrightarrow C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n)) \longrightarrow 0 \quad (1)$$

where

$$\begin{aligned} C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n)) : \quad & 0 \rightarrow \frac{R}{I^{n-d+1}} \rightarrow \cdots \rightarrow \left(\frac{R}{I^{n-d+i}}\right)^{\binom{d-1}{i}} \rightarrow \cdots \rightarrow \frac{R}{I^n} \rightarrow 0 \\ C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n)) : \quad & 0 \rightarrow \frac{R}{\mathfrak{m}I^{n-d+1}} \rightarrow \cdots \rightarrow \left(\frac{R}{\mathfrak{m}I^{n-d+i}}\right)^{\binom{d-1}{i}} \rightarrow \cdots \rightarrow \frac{R}{\mathfrak{m}I^n} \rightarrow 0 \\ C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n)) : \quad & 0 \rightarrow \frac{R}{I^{n-d+1}} \rightarrow \cdots \rightarrow \bigoplus_{j=0}^1 \left(\frac{R}{\mathfrak{m}^{1-j}I^{n-d+i-j}}\right)^{\binom{d-1}{i-j}} \rightarrow \cdots \rightarrow \frac{R}{\mathfrak{m}I^n} \rightarrow 0. \end{aligned}$$

Let $H_i(-)$ denote the i -th homology of the corresponding complex. From 1 we have the long exact sequence of homologies

$$\begin{aligned} \cdots & \rightarrow H_{i+1}(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) \rightarrow H_i(C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n))) \rightarrow \\ H_i(C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n))) & \rightarrow H_i(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) \rightarrow H_{i-1}(C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n))) \cdots \end{aligned} \quad (2)$$

We also have exact sequence:

$$0 \longrightarrow K.(\mathbf{a}_{d-1}^\circ, F(I))(n) \longrightarrow C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n)) \longrightarrow C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n)) \longrightarrow 0$$

gives rise to the long exact sequence:

$$\begin{aligned} \cdots & \rightarrow H_{i+1}(C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n))) \rightarrow H_{i+1}(C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n))) \rightarrow \\ H_i(K.(\mathbf{a}_{d-1}, F(I))(n)) & \rightarrow H_i(C(\mathbf{a}_{d-1}, \mathcal{F}, (1, n))) \rightarrow H_i(C(\mathbf{a}_{d-1}, \mathcal{F}, (0, n))) \end{aligned} \quad (3)$$

Lemma 3.5. *Let I be an $\mathfrak{m}$ -primary ideal of minimal mixed multiplicity in a Cohen-Macaulay local ring $(R, \mathfrak{m})$. Let $x \in \mathfrak{m}$ and $\mathbf{a}_{d-1} \in I$ be a joint reduction of $\mathfrak{m}$ and I . Then $H_i(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) = 0$ for all $n \geq 1$ and for all $i \geq 1$.*

Proof. It is enough to show that $H_1(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) = 0$ for all $n \geq 1$ [DG]. From [DG],

$$H_1(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) \cong \frac{(x, \mathbf{a}_{d-1}) \cap \mathfrak{m}I^n}{xI^n + \mathbf{a}_{d-1}\mathfrak{m}I^{n-1}}, \quad \text{for all } n \geq 1.$$

Since I is an ideal of minimal mixed multiplicity, for all $n \geq 1$, $\mathfrak{m}I^n = xI^n + \mathbf{a}_{d-1}^n\mathfrak{m}$. Hence

$$(x, \mathbf{a}_{d-1}) \cap \mathfrak{m}I^n \cong (x, \mathbf{a}_{d-1}) \cap (xI^n + \mathbf{a}_{d-1}^n\mathfrak{m}) = xI^n + \mathbf{a}_{d-1}^n\mathfrak{m} = \mathfrak{m}I^n.$$

Therefore $H_1(C(x, \mathbf{a}_{d-1}, \mathcal{F}, (1, n))) = 0$, for all $n \geq 1$. $\square$

Theorem 3.6. *Let I be an $\mathfrak{m}$ -primary ideal of minimal mixed multiplicity in a Cohen-Macaulay ring of dimension $d \geq 2$. Let $x, \mathbf{a}_{d-1}$ be a joint reduction of $\mathfrak{m}, I^{d-1}$. If $a_1^*, \dots, a_t^*$ is a regular sequence in $G(I)$, then $a_1^\circ, \dots, a_t^\circ \text{ depth}(F(I)) \geq d-1$. In particular, If $\text{depth}(G(I)) \geq d-1$, then $\text{depth}(F(I)) \geq d-1$.*

Proof. We can choose $x \in \mathfrak{m}$ and $a_1, \dots, a_{d-1} \in I$ to be a superficial sequence for $\mathfrak{m}$ and I , so that a_{d-1}^* is a superficial sequence in $G(I)$ and $F(I)$. Since $a_1^*, \dots, a_t^*$ is a regular sequence in $G(I)$ we have from [HM, Proposition 3.3], we get

$$H_i(C(\mathbf{a}_t, \mathcal{F}, (0, n))) = 0, \quad \text{for all } i \geq d - t + 1 \quad [\text{HM, Proposition 3.3}] \quad (4)$$

$$H_i(C(\mathbf{a}_t, \mathcal{F}, (1, n))), \quad \text{for all } i \geq d - t + 1, n \geq 0 \quad \text{from (2)} \quad (5)$$

$$H_i(K(\mathbf{a}_t, F(I))(n) = 0, \quad \text{for all } i \geq d - t + 1, n \geq 0 \quad \text{applying (4) and (5) to (3).} \quad (6)$$

□

Theorem 3.7. *Let I be an ideal of minimal mixed multiplicity in a Cohen-Macaulay local ring of dimension two. Let (x, a) be a joint reduction of I . If a° is a regular element in $F(I)$, then a^* is a regular element in $G(I)$. In particular if $\text{depth}(F(I)) \geq 1$, then $\text{depth}(G(I)) \geq 1$.*

Proof. Since I is an ideal of minimal mixed multiplicity, from Lemma 3.5, $H_i(C(x, a, \mathcal{F}, (1, n))) = 0$ for all $n \geq 1$ and for all $i \geq 1$. Hence from (2)

$$H_1(C(a, \mathcal{F}, (0, n))) \cong H_1(C(a, \mathcal{F}, (1, n))), \quad \text{for all } n \geq 0. \quad (7)$$

Since a° is a regular element in $F(I)$, from (3)

$$\begin{aligned} 0 &= H_0(C(a^\circ, F(I), n)) \\ H_1(C(a, \mathcal{F}, (0, n))) &\cong H_1(C(a, \mathcal{F}, (1, n))), \quad \text{for all } n \geq 0. \end{aligned} \quad (8)$$

and that the exact sequence:

$$0 \rightarrow H_0(C(a^\circ, F(I), n)) \rightarrow H_0(C(a, \mathcal{F}, (1, n))) \rightarrow H_0(C(a, \mathcal{F}, (0, n))) \rightarrow 0,$$

is exact for all $n \geq 0$, i.e.

$$0 \rightarrow \frac{I^n}{aI^{n-1} + \mathfrak{m}I^n} \rightarrow \frac{R}{a + \mathfrak{m}I^n} \rightarrow \frac{R}{a + I^n} \rightarrow 0, \quad \text{for all } n \geq 0.$$

Therefore

$$\frac{I^n}{aI^{n-1} + \mathfrak{m}I^n} \cong \frac{a + I^n}{a + \mathfrak{m}I^n} \cong \frac{I^n}{(a \cap I^n) + \mathfrak{m}I^n}.$$

This implies that

$$(a \cap I^n) \subseteq aI^{n-1} + \mathfrak{m}I^n = aI^{n-1} + xI^n.$$

The last inequality follows since I is an ideal of minimal mixed multiplicity. We claim that:

$$(a \cap I^n) = aI^{n-1} + ax(I^n : a).$$

Suppose $\alpha \in (a \cap I^n)$. Then $\alpha = a\beta = ai + xj$, where $\beta \in R$, $i \in I^{n-1}$ and $j \in I^n$. This implies that

$$a(\beta - i) = xj.$$

Since x, a is a regular sequence in R , $j = aj'$ where $j' \in (I^n : a)$. Therefore

$$(a \cap I^n) \subseteq aI^{n-1} + ax(I^n : a).$$

Clearly $aI^{n-1} + ax(I^n : a) \subseteq (a \cap I^n)$. Hence equality holds. This proves the claim.

Since $(a \cap I^n) = a(I^n : a)$ Hence we have

$$a(I^n : a) \subseteq aI^{n-1} + ax(I^n : a).$$

and by Nakayama's lemma,

$$a(I^n : a) = aI^{n-1}.$$

As a is a regular element in R ,

$$(I^n : a) = I^{n-1}, \quad \text{for all } n \geq 0.$$

Therefore a^* is a regular element in $G(I)$. Hence $\text{depth } G(I) \geq 1$. $\square$

Theorem 3.8. *Let I be a contracted ideal in a two dimensional regular local ring. Then $\text{depth}(R(I)) - 1 = \text{depth}(G(I)) = \text{depth}(F(I))$.*

Proof. By Theorem 2.2, $\text{depth}(R(I)) = \text{depth}(G(I)) - 1$. Hence we only need to prove that $\text{depth}(G(I)) = \text{depth}(F(I))$. By Corollary 2.5, $G(I)$ is Cohen-Macaulay if and only if $F(I)$ is Cohen-Macaulay.

By Theorem 2.5 and Theorem 3.7 we get $\text{depth}(G(I)) = 1$ if and only if $\text{depth}(F(I)) = 1$. The only case remaining is when $\text{depth}(G(I)) = 0$. In this case, $\text{depth}(F(I)) = 0$ and vice versa. $\square$

4. SOME EXAMPLES

We present an infinite class of examples where the Rees ring is Cohen-Macaulay but the fiber cone is not.

Example 4.1. Let $R = k[X, Y, Z]_{\mathfrak{m}}$, where $\mathfrak{m} = (X, Y, Z)$. Let $s \geq 2$ and $I_s = (Y^3, X^{s+1}Y^2, Y^2XZ, Y^2Z^3, X^{2s+1}Y, YZ^2X, YZ^5, X^{3s+1}, ZX^2, Z^4X, Z^7)$.

Then $J = (X^{3s+1} + Z^7, X^2Z, Y^3)$ is a minimal reduction of I . The following hold true:

- (i) $J \cap I^2 = JI$
- (ii) $JI^n = I^{n+1}$ for all $n \geq 2$.

Hence $R(I)$ and $G(I)$ is Cohen-Macaulay [GS], [HM].

It is easy to see that $\mu(I) := \ell(I/\mathfrak{m}I) = 11 = 9 + 2 = e(X + Y + Z, X^2Z, Y^3 + Z^7) + 2$. Hence I is an ideal of minimal mixed multiplicity and the Hilbert series of $F(I)$ is of the form [DRV, Theorem 2.4]

$$H(F(I), t) = \frac{1 + 8t}{(1 - t)^3}$$

Since I is an ideal of minimal mixed multiplicity and the reduction number of I , $r(I) > 1$, $F(I)$ is not Cohen-Macaulay [DRV, Corollary 2.5]. By Proposition 3.6, $\text{depth } F(I) = 2$. $\square$

We now present an infinite class of examples where the fiber cone is Cohen-Macaulay but the Rees ring is not.

Example 4.2. Let $R = k[X, Y]_{\mathfrak{m}}$, where $\mathfrak{m} = (X, Y)$. Let

$$I = (X^n, X^2Y^{n-2}, XY^{n-1}, Y^n).$$

Then $F(I) = k[X^n, X^2Y^{n-2}, XY^{n-1}, Y^n]$ is Cohen-Macaulay but $R(I)$ and $G(I)$ are not. Moreover if n is even then $F(I)$ is Gorenstein.

One can verify that

$$F(I) \cong \begin{cases} \frac{k[a, b, c, d]}{(c^2 - bc, b^{r+1} - acd^{r-1}, b^r c - ad^r)} & \text{if } n = 2r + 1 \\ \frac{k[a, b, c, d]}{(c^2 - bc, b^r - ad^{r-1})} & \text{if } n = 2r \end{cases}.$$

Clearly, for all $n \geq 4$, the image of a, d in $F(I)$ is a regular sequence. Hence $F(I)$ is Cohen-Macaulay.

Moreover, the Hilbert-Series of $F(I)$ is

$$H(F(I), t) = \begin{cases} \frac{1 + 2(t + \cdots + t^r)}{(1-t)^2} & \text{if } n = 2r + 1 \\ \frac{1 + 2(t + \cdots + t^{r-1}) + t^r}{(1-t)^2} & \text{if } n = 2r \end{cases}.$$

Put $J = (X^n, Y^n)$.

Put $r = \lfloor n/2 \rfloor$. Then $X^{n-1}Y^{n(r-1)+1} \in I^r \setminus JI^{r-1}$. and $JI^r = I^{r+1}$ which implies that the reduction number of I with respect to J is at least two. Now $J \cap I^2 \neq JI$ since $X^3Y^{2n-3} \in J \cap I^2 \setminus JI$. Therefore $G(I)$ is not Cohen-Macaulay. Also $R(I)$ are not Cohen-Macaulay. $\square$

REFERENCES

- [B] P. B. Bhattacharya, *The Hilbert function of two ideals*, Math. Proc. Cambridge Philos. Soc. **53** (1957), 568–575.
- [DG] C. D'Cruz; A. Guerrieri: *Koszul Homology, Mixed multiplicities and fiber cones* preprint.
- [AH] I. M. Aberbach; C. Huneke: *An improved Brianon-Skoda theorem with applications to the Cohen-Macaulayness of Rees algebras*. Math. Ann. **297** (1993), no. 2, 343–369.
- [CZ] T. Cortadellas; S. Zarzuela: *On the depth of the fiber cone of filtrations*. J. Algebra **198** (1997), no. 2, 428–445.
- [DRV] Clare D'Cruz, K. N. Raghavan and J. K. Verma, *Cohen-Macaulay fiber cones*, Commutative algebra, algebraic geometry and computational methods (Hanoi, 1996), Springer-Singapore (1999), 233–246.
- [DV] Clare D'Cruz; J. K. Verma, Hilbert series of fiber cones of ideals with almost minimal mixed multiplicity. J. Algebra 251 (2002), no. 1, 98–109.
- [GS] S. Goto, Shiro; Y. Shimoda: *On the Rees algebras of Cohen-Macaulay local rings*. Commutative algebra (Fairfax, Va., 1979), pp. 201–231, Lecture Notes in Pure and Appl. Math., **68**, Dekker, New York, 1982.
- [HKU] W. Heinzer, Mee-Kyoung Kim, B. Ulrich. *The Gorenstein and complete intersection properties of associated graded rings* preprint.

- [HM] S. Huckaba; T. Marley: *Depth properties of Rees algebras and associated graded rings*. J. Algebra **156** (1993), no. 1, 259–271.
- [J] A. V. Jayanthan: *On the depth of graded rings associated to lex-segment ideals in $k[x, y]$* , Beitrge Algebra Geom. (to appear).
- [Sh] K. Shah, Kishor *On the Cohen-Macaulayness of the fiber cone of an ideal*. J. Algebra **143** (1991), no. 1, 156–172.
- [V] J. Verma; *Rees algebras and mixed multiplicities*. Proc. Amer. Math. Soc. **104** (1988), no. 4, 1036–1044.
- [ZS] O. Zariski, P. Samuel: “*Commutative Algebra*”, Vol II, Van Nostrand, Princeton, NJ, 1960.

CHENNAI MATHEMATICAL INSTITUTE, 92, G.N. CHETTY ROAD, CHENNAI 600 017, INDIA.

E-mail address: `clare@cmi.ac.in`